


Telkom
University



ELEKTROMAGNETIK TERAPAN

1. VEKTOR ANALISIS



By Dwi Andi Nurmantris

WHERE ARE WE?

- 1. PENDAHULUAN**
 - Pengenalan Mata Kuliah
 - Silabus (materi), dan referensi,
 - Aturan penilaian: Quis, Ujian, Tugas dll
 - Aturan Perkuliahahan : Kontrak Belajar
 - Sejarah dan Aplikasi Elektromagnetika
- 2. PERSAMAAN MAXWELL UNTUK MEDAN DINAMIS**
 - Vektor Analysis
 - Konsep dan Arti Fisis tentang Empat Persamaan Maxwell
 - penerapan konsep Empat Persamaan Maxwell
- 3. PROPAGASI GELOMBANG DATAR**
 - Penurunan Pers. Helmholtz dari Persamaan Maxwell
 - Perambatan gelombang pada Berbagai Medium (Dielektrik Merugi)
 - Perambatan gelombang pada Dielektrik Sempurna, Vakum, Sekundernya
 - Vektor Poynting dan Analisis Daya
 - Polarisasi Gelombang
 - Pantulan gelombang sudut datang nol
 - Konservasi Daya dalam Pantulan
 - Standing Wave Ratio, Impedansi Input, dan Matching gelombang
 - Radome (med1|med2|med3 -med1|med2|med3)
 - Perambatan GEM pada arah sembarang
 - Pantulan Sudut-Datang Tak-Nol dan Nol : Gelombang Berdiri
- 4. SALURAN TRANSMISI**
 - Model dan Persamaan Saluran Transmisi
 - Macam-macam Saluran Transmisi dengan Parameter Primer dan Sekundernya, Saluran Distortionless dan Lossless

WHERE ARE WE?

- Kasus 1 : Saluran Tak-merugi Beban Sesuai (V , I , P)
- Kasus 2 : Saluran Tak-merugi Beban Tak-Sesuai (V , I , P)
- Impedansi input dan VSWR
- Kasus 3 : Saluran-saluran Istimewa ($\lambda/2$, $\lambda/4$, $Z_L = 0$, $Z_L = \infty$)
- Kasus 4 : Persoalan Saluran Merugi
- Penyesuaian Impedansi dengan Transformator $1/4$ panjang glb.
- Konsep lebar-pita frekuensi untuk sistem saluran transmisi
- Penyesuaian Impedansi dengan Stub-Tunggal
- Smith-Chart: Pembuatan dan Penggunaan
- Penyesuaian Impedansi dengan Stub Ganda dengan Smith Chart

5. BUMBUNG GELOMBANG PERSEGI (BGP)

- Analisis Medan Elektromagnetik dalam BGP
- Gelombang Mode TM_{mn}, Parameter Primer dan Sekunder
- Gelombang Mode TE_{mn}, Parameter Primer dan Sekunder
- Tinjauan Daya dan Rugi-rugi

6. BUMBUNG GELOMBANG SIRKULAR (BGS)


- Analisis Medan Elektromagnetik dalam BGS
- Gelombang Mode TM_{nl} dan TE_{nl} , Parameter Primer dan Sekunder
- Pengenalan Serat Optik

7. RADIASI GELOMBANG


- Analisis Medan Radiasi Filamen Pendek, Diagram Arah
- Aproksimasi untuk Medan Jauh, Daya Pancar, Tahanan Pancar
- Dipole $1/2\lambda$ dan Monopole

OUTLINE



- 
- The slide features a decorative border composed of red and green wavy lines that frame the central content area.
1. Vektor Analysis
 - a) Scalar dan vektor
 - b) Vektor Algebra
 - c) Coordinate system
 - d) Calculus of scalar and vektor
(gradient, divergence, curl)
 - e) Divergence Theorem
 - f) Stoke's Theorem

MAXWELL'S EQUATION



MAXWELL'S EQUATION



This is where we are heading

$$\nabla \cdot \vec{D} = \rho$$

$$\nabla \cdot \vec{B} = 0$$

$$\nabla \times \vec{E} = - \frac{\partial \vec{B}}{\partial t}$$


$$\nabla \times \vec{H} = \sigma \vec{E} + \frac{\partial \vec{D}}{\partial t}$$

$$\oint_s \vec{D} \cdot d\vec{s} = \int_v \rho \, dv$$

$$\oint_s \vec{B} \cdot d\vec{s} = 0$$

$$\oint_s \vec{E} \cdot d\vec{l} = - \oint_s \frac{\partial \vec{B}}{\partial t} \cdot d\vec{s}$$

$$\int_l \vec{H} \cdot d\vec{l} = \oint_s (\sigma \vec{E} + \frac{\partial \vec{D}}{\partial t}) \cdot d\vec{s}$$



James Clark Maxwell.



DON'T
PANIC!

WHY MAXWELL'S EQUATION?



- How EM field produced?
- How EM wave produced?
- How does a capacitor store energy?
- How does an antenna radiate or receive signals?
- How do Electromagnetic fields propagate in space?
- What really happens when electromagnetic energy travels from one end of waveguide to the others?

OUTLINE



1. Vektor Analysis

- a) Scalar and vector
- b) Vektor Algebra
- c) Coordinate system
- d) Calculus of scalar and vektor

Scalar and Vector

Skalar : Besaran yang hanya memiliki nilai

Contoh : Temperatur, laju, jarak, dll

- a. **Scalar Discrete Quantities** : can be described with a single numeric value
- b. **Scalar Fields Quantities** : that the magnitude must be described as one Function of (typically) space and/or time

Vektor : Besaran yang memiliki nilai dan arah

Contoh : Medan listrik, medan magnet, Gaya, kecepatan, posisi, percepatan, dll

- a. **Vector Discrete Quantities** : must be described with both a single value magnitude and a direction
- b. **Vector Fields Quantities**: quantities that both magnitude and/or direction must be described as a function of (typically) space and/or time

Scalar and Vector

Contoh Physical Quantity dalam scalar dan vektor

- **Scalar Quantities**

- ✓ Berat
- ✓ Tinggi
- ✓ Temperatur

- **Vector Quantities**

- ✓ Force
- ✓ velocities

- **Scalar Fields**


- ✓ Berat dalam fungsi waktu
- ✓ Surface Temperatur saat ini pada area tertentu

- **Vector Fields**

- ✓ Surface wind velocities



Vector Representation

symbolically we can represent a vector quantity as an arrow:



- ❑ The length of the arrow is proportional to the magnitude of the vector quantity.
- ❑ The orientation of the arrow indicates the direction of the vector quantity.
- ❑ The variable names of a vector quantity will always be either boldface or have an overbar

Dua buah vektor dikatakan sama jika kedua vektor tersebut memiliki magnitude dan arah yang identik



Vector Representation

$$\vec{A} = |\vec{A}| \hat{a}_A$$

Dengan :

$|\vec{A}|$ adalah besar vektor A atau panjang vektor A (magnitude vektor A)

\hat{a}_A adalah unit vektor A atau vektor satuan searah A

Vektor satuan atau unit vektor menyatakan arah vektor, besarnya satu.




$$\hat{a}_A = \frac{\vec{A}}{|\vec{A}|}$$

$$|\hat{a}_A| = 1$$


Vector Algebra

If we know the rules of vector operations, we can analyze, manipulate, and simplify vector operations!




Vector Addition


$$\vec{A} + \vec{B} = \vec{C}$$



1. Vector addition is commutative $\rightarrow \vec{A} + \vec{B} = \vec{B} + \vec{A}$





2. Vector addition is associative $\rightarrow (\vec{X} + \vec{Y}) + \vec{Z} = \vec{X} + (\vec{Y} + \vec{Z})$



Vector Algebra



Head-to-tail Vektor Composition





$$\vec{f} = \vec{a} + \vec{b} + \vec{c} + \vec{d} + \vec{e}$$


Vector Algebra

Vector Subtraction

the negative of a vector is a vector with equal magnitude but opposite direction.



$$A + (-B) = A - B$$




Vector Algebra

Scalar-Vector multiplication


the product of a scalar and a vector—is a vector!

$$a\vec{B} = \vec{C} = a|\vec{B}|\hat{a}_B$$




1. scalar-vector multiplication is distributive:

$$a B + b B = (a+b) B$$



$$a B + a C = a (B+C)$$



2. scalar-vector multiplication is Commutative:

$$aB = B a$$

3. Multiplication of a vector by a negative scalar:

$$-a B = a(-B)$$



4. Division of a vector by a scalar is the same as multiplying the vector by the inverse of the scalar


$$\frac{\vec{B}}{a} = \left(\frac{1}{a}\right)\vec{B}$$

Vector Algebra

Dot Product / Perkalian Skalar

The dot product of two vectors is defined as:

$$\mathbf{A} \cdot \mathbf{B} = |\mathbf{A}| |\mathbf{B}| \cos \theta_{AB}$$



IMPORTANT NOTE: The dot product is an operation involving two vectors, but the result is a scalar

1. the dot product is commutative

$$\vec{A} \bullet \vec{B} = \vec{B} \bullet \vec{A}$$

2. The dot product is distributive with addition


$$\vec{A} \bullet (\vec{B} + \vec{C}) = \vec{A} \bullet \vec{B} + \vec{A} \bullet \vec{C}$$

A.B.C = ???



3. $\vec{A} \bullet \vec{A} = |\vec{A}| |\vec{A}| \cos 0^\circ = |\vec{A}|^2$

4.



$$\vec{A} \bullet \vec{B} = 0$$

5



$$\vec{A} \bullet \vec{B} = |\vec{A}| |\vec{B}| \cos 0^\circ = |\vec{A}| |\vec{B}|$$

6.




$$\vec{A} \bullet \vec{B} = |\vec{A}| |\vec{B}| \cos 180^\circ = -|\vec{A}| |\vec{B}|$$

Vector Algebra


Cross Product

the product of a scalar and a vector—is a vector!


$$\mathbf{A} \times \mathbf{B} = \hat{a}_n |\mathbf{A}| |\mathbf{B}| \sin \theta_{AB}$$



IMPORTANT NOTE: The cross product is an operation involving two vectors, and the result is also a vector



1.



$$\vec{A} \times \vec{B} = \hat{a}_n |\vec{A}| |\vec{B}| \sin 90^\circ = \hat{a}_n |\vec{A}| |\vec{B}|$$

2.



$$\vec{A} \times \vec{B} = \hat{a}_n |\vec{A}| |\vec{B}| \sin 0^\circ = \hat{a}_n |\vec{A}| |\vec{B}| \sin 180^\circ = 0$$

3. The cross product is not commutative

$$\vec{A} \times \vec{B} \neq \vec{B} \times \vec{A}$$

4. The cross product is also not associative

$$(\vec{A} \times \vec{B}) \times \vec{C} \neq \vec{A} \times (\vec{B} \times \vec{C})$$

5. the cross product is distributive

$$\vec{A} \times (\vec{B} + \vec{C}) = (\vec{A} \times \vec{B}) + (\vec{A} \times \vec{C})$$

Vector Algebra

Triple Product

The triple product is simply a combination of the dot and cross products.


$$\vec{A} \bullet \vec{B} \times \vec{C} = \vec{A} \bullet (\vec{B} \times \vec{C})$$

IMPORTANT NOTE: The triple product $\vec{A} \bullet \vec{B} \times \vec{C}$ results in a scalar value.

The Cyclic Property

$$\vec{A} \bullet \vec{B} \times \vec{C} = \vec{C} \bullet \vec{A} \times \vec{B} = \vec{B} \bullet \vec{C} \times \vec{A}$$

The cyclical rule means that the triple product is invariant to Shifts in the order of the vectors



Vector Algebra



Let's test your vector algebraic skills!
Can you evaluate the following expressions,
and determine whether the result is a
scalar (S), a vector(V), or neither (N) ??

IMPORTANT NOTE: If the expression initially results in a vector (or scalar), then after each manipulation, the result must also be a vector (or scalar)

1. $(\mathbf{A} \cdot \mathbf{B})\mathbf{C}$ _____

2. $\mathbf{A} + (\mathbf{B} \cdot \mathbf{C})$ _____

3. $\mathbf{A} \cdot (\mathbf{B} \cdot \mathbf{C})$ _____

4. $\mathbf{A}(\mathbf{B} \times \mathbf{C})$ _____

5. $\mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$ _____

6. $\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) + \mathbf{C} \cdot (\mathbf{A} + \mathbf{B})$ _____

7. $\mathbf{A} \cdot \mathbf{B} \times \mathbf{C} \cdot \mathbf{D}$ _____

Intermezo




Find the baby

Coordinate Systems


Coordinates system

- A set of 3 scalar values that define position and a set of 3 unit vectors that define direction form a Coordinate system
- The 3 scalar values used to define position are called coordinates
- All coordinates are defined with respect to an arbitrary point called the origin.
- The 3 unit vectors used to define direction are called base vectors


Cartesian Coordinate



Cylindrical Coordinate




Spherical Coordinate



Coordinate Systems



Cartesian Coordinates system



- Kita dapat menentukan posisi (koordinat) titik P dengan 3 scalar values yaitu x, y, dan z
- coordinate values in the Cartesian system effectively represent the distance from a plane intersecting the origin
 - to specify the direction of a vector quantity is by using a well defined orthonormal set of vectors known as base vectors. \hat{a}_x , \hat{a}_y , \hat{a}_z
- Orthonormal :
 - a. Each vector is a unit vector:
$$\hat{a}_x \bullet \hat{a}_x = \hat{a}_y \bullet \hat{a}_y = \hat{a}_z \bullet \hat{a}_z = 1$$
 - a. Each vector is mutually orthogonal:
$$\hat{a}_x \bullet \hat{a}_y = \hat{a}_y \bullet \hat{a}_z = \hat{a}_x \bullet \hat{a}_z = 0$$
- Additionally, a set of base vectors must be arranged such that:
$$\hat{a}_x \times \hat{a}_y = \hat{a}_z; \hat{a}_y \times \hat{a}_z = \hat{a}_x; \hat{a}_z \times \hat{a}_x = \hat{a}_y$$

Coordinate Systems

Cylindrical Coordinates system




- Kita dapat menentukan posisi (koordinat) titik P pada koordinat tabung dengan 3 scalar values yaitu ρ, φ, z
- Keterangan :
 - 1) The value ρ indicates the distance of the point from the z -axis ($0 \leq \rho < \infty$)
 - 2) The value φ indicates the rotation angle around the z -axis ($0 \leq \varphi < 2\pi$)
 - 3) The value z indicates the distance of the point from the $x-y$ ($z = 0$) plane ($-8 < z < 8$)
- to specify the direction of a vector quantity is by using a base vectors \hat{a}_ρ \hat{a}_φ \hat{a}_z
- Not like Cartesian base vectors, the cylindrical base vectors are dependent on position (\hat{a}_ρ \hat{a}_φ)
- set of base vectors must be arranged such that:

$$\hat{a}_\varphi \times \hat{a}_z = \hat{a}_\rho; \quad \hat{a}_z \times \hat{a}_\rho = \hat{a}_\varphi; \quad \hat{a}_\rho \times \hat{a}_\varphi = \hat{a}_z$$

Coordinate Systems

Spherical Coordinates system




- Kita dapat menentukan posisi (koordinat) titik P pada koordinat tabung dengan 3 scalar values yaitu r, θ, φ
- Keterangan :
 - 1) The value r ($0 = r < \infty$) expresses the distance of the point from the origin.
 - 2) Angle θ ($0 \leq \theta \leq \pi$) represents the angle formed with the z-axis
 - 3) Angle φ ($0 \leq \varphi < 2\pi$) represents the rotation angle around the z-axis
- to specify the direction of a vector quantity is by using a base vectors $\hat{a}_r, \hat{a}_\varphi, \hat{a}_\theta$
- the Spherical base vectors are dependent on position ($\hat{a}_r, \hat{a}_\varphi, \hat{a}_\theta$)
- set of base vectors must be arranged such that:

$$\hat{a}_\theta \times \hat{a}_\varphi = \hat{a}_r; \quad \hat{a}_\varphi \times \hat{a}_r = \hat{a}_\theta; \quad \hat{a}_r \times \hat{a}_\theta = \hat{a}_\varphi$$


Coordinate Systems

1. Posisi/coordinate suatu titik pada sistem koordinat

Titik A berposisi pada $x=2$, $y=3$, dan $z=4$ atau $A(2, 3, 4)$ pada koordinat kertesian



Titik B berposisi pada $r=2$, $\theta=90^\circ$, dan $z=4$ atau $B(2, 3, 4)$ pada koordinat tabung




Coba gambar koordinat titik C pada koordinat bola dimana $r=4$, $\theta=90^\circ$, $\phi=45^\circ$



Coordinate Systems

2. Position Vector pada sistem koordinat

a vector beginning at the origin and extending outward to a point—is a very important and fundamental directed distance known as the position vector \bar{r}



The magnitude of \bar{r}

$$|\bar{r}| = \sqrt{\bar{r} \cdot \bar{r}} = \sqrt{x^2 + y^2 + z^2} = r$$

Vektor satuan searah vektor \bar{r}

$$\hat{a}_r = \frac{\bar{r}}{|\bar{r}|} = \frac{x\hat{a}_x + y\hat{a}_y + z\hat{a}_z}{\sqrt{x^2 + y^2 + z^2}}$$

Using the Cartesian coordinate system, the position vector can be explicitly written as:

$$\bar{r} = x \hat{a}_x + y \hat{a}_y + z \hat{a}_z$$

Contoh : $\bar{r} = \hat{a}_x + 2\hat{a}_y - 3\hat{a}_z$

Sehingga dengan mudah kita bisa langsung menentukan koordinat titik yang dituju. X=1, y=2, z=-3

NEVER, EVER express the position vector in either of Cylindrical base vector or Spherical base vector

$$\bar{r} \neq \rho \hat{a}_\rho + \phi \hat{a}_\phi + z \hat{a}_z$$

$$\bar{r} \neq r \hat{a}_r + \theta \hat{a}_\theta + \phi \hat{a}_\phi$$

Coordinate Systems


Latihan Soal

Titik A(3,4,5) terletak dalam koordinat Cartesius. Tentukan :

- a) Gambar vektor posisi A
- b) Tulis vektor posisi A
- c) Hitung besar/magnitude vektor A
- d) Tentukan vektor satuan searah A

Jarak antar 2 titik (koordinat)

- ❑ Position Vector berguna saat kita ingin menentukan jarak antar 2 titik dalam suatu sistem koordinat
- ❑ Misalkan dua buah titik P dan Q



$$\vec{P} = p_x \hat{a}_x + p_y \hat{a}_y + p_z \hat{a}_z$$

$$\vec{Q} = q_x \hat{a}_x + q_y \hat{a}_y + q_z \hat{a}_z$$

- ❑ Jarak antara dua titik P dan Q adalah magnitudo dari perbedaan vektor P dan Q

$$\overrightarrow{PQ} = (q_x - p_x) \hat{a}_x + (q_y - p_y) \hat{a}_y + (q_z - p_z) \hat{a}_z$$

- ❑ Jarak titik P dan Q

$$D_{PQ} = \sqrt{(q_x - p_x)^2 + (q_y - p_y)^2 + (q_z - p_z)^2}$$

Coordinate Systems

3. Scalar Field pada sistem koordinat

- a scalar field is a scalar quantity that is a function of position and/or time
- Scalar field can be written as :

$$A(x, y, z) = f(x, y, z)$$

$$A(\rho, \varphi, z) = f(\rho, \varphi, z)$$

$$A(r, \theta, \varphi) = f(r, \theta, \varphi)$$

Contoh

$$g(x, y, z) = (x^2 + y^2)yz$$

Coba Anda Gambarkan
scalar field disamping
pada koordinat kartesian



Coordinate Systems

4. Vector pada sistem koordinat

any vector can be written as a sum of three vectors!

$$\vec{A} = A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z$$

$$\vec{A} = A_\rho \hat{a}_\rho + A_\phi \hat{a}_\phi + A_z \hat{a}_z$$

$$\vec{A} = A_r \hat{a}_r + A_\phi \hat{a}_\phi + A_\theta \hat{a}_\theta$$

- Each of these three vectors point in one of the three orthogonal directions
- The magnitude of each of these three vectors are determined by the scalar values A_x , A_y , and A_z that called the scalar components of vector A.
- The vectors $A_x \hat{a}_x$, $A_y \hat{a}_y$, $A_z \hat{a}_z$ are called the vector components of A.

- Menentukan scalar component :

$$\begin{aligned}\vec{A} \bullet \hat{a}_x &= (A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z) \bullet \hat{a}_x \\ &= A_x \hat{a}_x \bullet \hat{a}_x + A_y \hat{a}_y \bullet \hat{a}_x + A_z \hat{a}_z \bullet \hat{a}_x = A_x\end{aligned}$$

$A_x = \vec{A} \bullet \hat{a}_x$

$A_y = \vec{A} \bullet \hat{a}_y$

$A_z = \vec{A} \bullet \hat{a}_z$

- Sehingga :

$$\vec{A} = (\vec{A} \bullet \hat{a}_x) \hat{a}_x + (\vec{A} \bullet \hat{a}_y) \hat{a}_y + (\vec{A} \bullet \hat{a}_z) \hat{a}_z$$

- Menentukan Magnitude Vector:


$$\begin{aligned}|\vec{A}| &= \sqrt{A_x^2 + A_y^2 + A_z^2} \\ &= \sqrt{A_\rho^2 + A_\phi^2 + A_z^2} \\ &= \sqrt{A_r^2 + A_\theta^2 + A_\phi^2}\end{aligned}$$

- Menentukan vektor satuan searah vektor A

$$\hat{a}_A = \frac{\vec{A}}{|\vec{A}|} = \frac{A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z}{\sqrt{A_x^2 + A_y^2 + A_z^2}}$$

Coordinate Systems

Contoh



$$\begin{aligned} A_x &= \vec{A} \bullet \hat{a}_x = 2 & A_r &= \vec{A} \bullet \hat{a}_r = 0 \\ A_y &= \vec{A} \bullet \hat{a}_y = 1,5 & A_\varphi &= \vec{A} \bullet \hat{a}_\varphi = 2,5 \\ A_z &= \vec{A} \bullet \hat{a}_z = 0 & A_\theta &= \vec{A} \bullet \hat{a}_\theta = 0 \end{aligned}$$

$$\overbrace{\quad\quad\quad}^{\text{Curly brace}}$$



$$\vec{A} = A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z$$

$$= 2 \hat{a}_x + 1,5 \hat{a}_y$$

$$\vec{A} = 2,5 \hat{a}_\varphi$$

$$\left. \begin{aligned} |\vec{A}| &= \sqrt{A_x^2 + A_y^2 + A_z^2} \\ &= \sqrt{2^2 + 1,5^2} = \sqrt{6,25} = 2,5 \\ &= \sqrt{A_r^2 + A_\theta^2 + A_\varphi^2} \quad \hat{a}_A = \frac{\vec{A}}{|\vec{A}|} = \frac{2 \hat{a}_x + 1,5 \hat{a}_y}{2,5} = 0,8 \hat{a}_x + 0,6 \hat{a}_y \\ &= \sqrt{2,5^2} = 2,5 \end{aligned} \right\}$$

$$\hat{a}_A = \frac{\vec{A}}{|\vec{A}|} = \frac{2,5 \hat{a}_\varphi}{2,5} = \hat{a}_\varphi$$



Coordinate Systems

Latihan soal

1. Gambarkan vektor berikut dalam sistem koordinat silinder
 - a) $A = 3a_\rho + 2a_\phi + a_z$ berpangkal di $M(2,0,0)$
 - b) $B = 3a_\rho + 2a_\phi + a_z$ berpangkal di $N(2,\pi/2,0)$
2. Gambarkan vektor berikut pada sistem koordinat bola
 - a) $C = 3a_r + a_\theta + 2a_\phi$ berpangkal di $M(2, \pi/2, 0)$
 - b) $D = 3a_r + a_\theta + 2a_\phi$ berpangkal di $N(2, \pi/2, \pi/2)$

Coordinate Systems

5. Vector Fields pada sistem koordinat

- ❑ a vector field is a vector quantity that is a function of position and/or time
- ❑ When we express a vector field using orthonormal base vectors, the scalar component of each direction is a scalar field—a scalar function of position!

$$\left. \begin{array}{l} \vec{A} = A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z \\ \vec{A} = A_\rho \hat{a}_\rho + A_\varphi \hat{a}_\varphi + A_z \hat{a}_z \\ \vec{A} = A_r \hat{a}_r + A_\varphi \hat{a}_\varphi + A_\theta \hat{a}_\theta \end{array} \right\} \begin{array}{l} \vec{A}(x, y, z) = A_x(x, y, z) \hat{a}_x + A_y(x, y, z) \hat{a}_y + A_z(x, y, z) \hat{a}_z \\ \vec{A}(\rho, \varphi, z) = A_\rho(\rho, \varphi, z) \hat{a}_\rho + A_\varphi(\rho, \varphi, z) \hat{a}_\varphi + A_z(\rho, \varphi, z) \hat{a}_z \\ \vec{A}(r, \varphi, \theta) = A_r(r, \varphi, \theta) \hat{a}_r + A_\varphi(r, \varphi, \theta) \hat{a}_\varphi + A_\theta(r, \varphi, \theta) \hat{a}_\theta \end{array}$$

Contoh

$$\vec{A}(x, y, z) = (x^2 + y^2) \hat{a}_x + \frac{xz}{y} \hat{a}_y + (3 - y) \hat{a}_z$$

- ❑ Often, vector fields must be written as variables of three spatial coordinates, as well as a time variable t and known as a dynamic vector field.

Contoh


$$\vec{A}(x, y, z, t) = (x^2 + y^2) t \hat{a}_x + \frac{xz}{y} t^2 \hat{a}_y + (3 - y + 4t) \hat{a}_z$$




Coordinate Systems

Gallery of Vector Scalar and Vector Field


$$\vec{A} = \hat{a}_x$$



$$\vec{A} = x\hat{a}_x + y\hat{a}_y$$



$$\vec{A} = x\hat{a}_x$$



Coba gambarkan vector field beikut :


$$\vec{A} = \hat{a}_\rho$$



Coordinate Systems

Point Location/Coordinate and scalar field Transformation

Transformation of expressing point location/Coordinate and scalar field in terms of cartesian coordinate to cylindrical or spherical coordinates or vice versa



Cartesian ke Cylindrical

$$(x, y, z) \Rightarrow (\rho, \varphi, z)$$

$$\rho = \sqrt{x^2 + y^2}$$

$$\varphi = \tan^{-1}\left(\frac{y}{x}\right)$$

$$z = z$$

Cylindrical ke Cartesian

$$(\rho, \varphi, z) \Rightarrow (x, y, z)$$

$$x = \rho \cdot \cos \varphi$$

$$y = \rho \cdot \sin \varphi$$

$$z = z$$

Cartesian ke Spherical

$$(x, y, z) \Rightarrow (r, \theta, \varphi)$$

$$r = \sqrt{x^2 + y^2 + z^2}$$

$$\varphi = \tan^{-1}\left(\frac{y}{x}\right)$$

$$\theta = \tan^{-1}\left(\frac{\sqrt{x^2 + y^2}}{z}\right)$$

Spherical ke Cartesian

$$(r, \theta, \varphi) \Rightarrow (x, y, z)$$

$$x = r \cdot \sin \theta \cdot \cos \varphi$$

$$y = r \cdot \sin \theta \cdot \sin \varphi$$

$$z = r \cdot \cos \theta$$

Cylindrical ke Spherical

$$(\rho, \varphi, z) \Rightarrow (r, \theta, \varphi)$$

$$r = \sqrt{\rho^2 + z^2}$$

$$\theta = \tan^{-1}\left(\frac{\rho}{z}\right)$$

$$\varphi = \varphi$$

Spherical ke Cylindrical

$$(r, \theta, \varphi) \Rightarrow (\rho, \varphi, z)$$

$$\rho = r \sin \theta$$

$$\varphi = \varphi$$

$$z = r \cos \theta$$



Coordinate Systems

Latihan Soal

1. Titik A(-3,-3,2) terletak dalam koordinat Cartesius. Tentukan :
 - a. Tentukan koordinat titik tersebut pada koordinat tabung
 - b. Tentukan koordinat titik tersebut pada koordinat bola
2. Suatu scalar field $g(\rho, \varphi, z) = \rho^3 z \sin \varphi$
tulis fungsi g di atas dengan expresi koordinat kartesian

Coordinate Systems

Vector Transformation

Transformation of expressing Vector field in terms of cartesian coordinate to cylindrical or spherical coordinates or vice versa

Vektor Field di Koordinat Cartesian

$$\mathbf{A}(x, y, z) = A_x(x, y, z) \hat{a}_x + A_y(x, y, z) \hat{a}_y + A_z(x, y, z) \hat{a}_z$$



$$\mathbf{A} = (\mathbf{A} \cdot \hat{a}_x) \hat{a}_x + (\mathbf{A} \cdot \hat{a}_y) \hat{a}_y + (\mathbf{A} \cdot \hat{a}_z) \hat{a}_z$$

| | | | |
|-------------|----------------|----------------|-------------|
| • | \vec{a}_ρ | \vec{a}_ϕ | \vec{a}_z |
| \vec{a}_x | $\cos \phi$ | $-\sin \phi$ | 0 |
| \vec{a}_y | $\sin \phi$ | $\cos \phi$ | 0 |
| \vec{a}_z | 0 | 0 | 1 |

| | | | |
|-------------|-----------------------|-----------------------|----------------|
| . | \vec{a}_r | \vec{a}_θ | \vec{a}_ϕ |
| \vec{a}_x | $\sin\theta \cos\phi$ | $\cos\theta \cos\phi$ | $-\sin\phi$ |
| \vec{a}_y | $\sin\theta \sin\phi$ | $\cos\theta \sin\phi$ | $\cos\phi$ |
| \vec{a}_z | $\cos\theta$ | $-\sin\theta$ | 0 |

Vektor Field di Koordinat Silinder


$$\begin{aligned}\vec{A}(\rho, \varphi, z) &= A_\rho(\rho, \varphi, z) \hat{a}_\rho + A_\varphi(\rho, \varphi, z) \hat{a}_\varphi + A_z(\rho, \varphi, z) \hat{a}_z \\ &= (\vec{A}(x, y, z) \bullet \hat{a}_\rho) \hat{a}_\rho + (\vec{A}(x, y, z) \bullet \hat{a}_\varphi) \hat{a}_\varphi + (\vec{A}(x, y, z) \bullet \hat{a}_z) \hat{a}_z\end{aligned}$$

Vektor Field di Koordinat Bola

$$\begin{aligned}\vec{A}(r, \theta, \varphi) &= A_r(r, \theta, \varphi) \hat{a}_r + A_\theta(r, \theta, \varphi) \hat{a}_\theta + A_\varphi(r, \theta, \varphi) \hat{a}_\varphi \\ &= (\vec{A}(x, y, z) \bullet \hat{a}_r) \hat{a}_r + (\vec{A}(x, y, z) \bullet \hat{a}_\theta) \hat{a}_\theta + (\vec{A}(x, y, z) \bullet \hat{a}_\varphi) \hat{a}_\varphi\end{aligned}$$

Coordinate Systems

Vector Transformation



$$\vec{A}(x, y, z) = A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z$$

$$\vec{A}(\rho, \phi, z) = A_\rho \hat{a}_\rho + A_\phi \hat{a}_\phi + A_z \hat{a}_z$$

$$\vec{A}(r, \theta, \phi) = A_r \hat{a}_r + A_\theta \hat{a}_\theta + A_\phi \hat{a}_\phi$$

Cartesian ke Cylindrical

$$(x, y, z) \Rightarrow (\rho, \phi, z)$$

$$A_\rho = A_x \cos \phi + A_y \sin \phi$$

$$A_\phi = -A_x \sin \phi + A_y \cos \phi$$

$$A_z = A_z$$

Cylindrical ke Cartesian

$$(\rho, \phi, z) \Rightarrow (x, y, z)$$

$$A_x = A_\rho \cos \phi - A_\phi \sin \phi$$

$$A_y = A_\rho \sin \phi + A_\phi \cos \phi$$

$$A_z = A_z$$



Cartesian ke Spherical

$$(x, y, z) \Rightarrow (r, \theta, \phi)$$

$$A_r = A_x \cos \phi \sin \theta + A_y \sin \phi \sin \theta + A_z \cos \theta$$

$$A_\theta = A_x \cos \phi \cos \theta + A_y \sin \phi \cos \theta - A_z \sin \theta$$

$$A_\phi = -A_x \sin \phi + A_y \cos \phi$$

Spherical ke Cartesian

$$(r, \theta, \phi) \Rightarrow (x, y, z)$$

$$A_x = A_r \sin \theta \cos \phi + A_\theta \cos \theta \cos \phi - A_\phi \sin \phi$$

$$A_y = A_r \sin \theta \sin \phi + A_\theta \cos \theta \sin \phi + A_\phi \cos \phi$$

$$A_z = A_r \cos \theta - A_\theta \sin \theta$$

Cylindrical ke Spherical

$$(\rho, \phi, z) \Rightarrow (r, \theta, \phi)$$

$$A_r = A_\rho \sin \theta + A_z \cos \theta$$

$$A_\theta = A_\rho \cos \theta - A_z \sin \theta$$

$$A_\phi = A_\phi$$

Spherical ke Cylindrical


$$(r, \theta, \phi) \Rightarrow (\rho, \phi, z)$$

$$A_\rho = A_r \sin \theta + A_\theta \cos \theta$$

$$A_\phi = A_\phi$$

$$A_z = A_r \cos \theta - A_\theta \sin \theta$$

Coordinate Systems



Latihan soal

1. Vektor $\mathbf{A} = 3\hat{\mathbf{a}}_\rho + 4\hat{\mathbf{a}}_\phi + 5\hat{\mathbf{a}}_z$ berada pada sistem koordinat silinder dengan titik pangkal di $(10, \pi/2, 0)$. Tentukan penulisan vektor ini pada sistem koordinat kartesian!
2. Suatu vektor $\vec{B}(x, y, z) = y\hat{\mathbf{a}}_x - x\hat{\mathbf{a}}_y + z\hat{\mathbf{a}}_z$ tentukan:
 - a. Expresi vektor tersebut pada koordinat tabung!
 - b. Expresi vektor tersebut pada koordinat bola!

Coordinate Systems

Vektor Algebra pada coordinate systems

Misalnya kita memiliki dua buah vektor pada koordinat kartesian :

$$\mathbf{A} = A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z$$

$$\mathbf{B} = B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z$$

Addition and Subtraction

$$\begin{aligned}\mathbf{A} + \mathbf{B} &= (A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z) + (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x \hat{a}_x + B_x \hat{a}_x + A_y \hat{a}_y + B_y \hat{a}_y + A_z \hat{a}_z + B_z \hat{a}_z \\ &= (A_x + B_x) \hat{a}_x + (A_y + B_y) \hat{a}_y + (A_z + B_z) \hat{a}_z\end{aligned}$$

$$\begin{aligned}\mathbf{A} - \mathbf{B} &= (A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z) - (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x \hat{a}_x - B_x \hat{a}_x + A_y \hat{a}_y - B_y \hat{a}_y + A_z \hat{a}_z - B_z \hat{a}_z \\ &= (A_x - B_x) \hat{a}_x + (A_y - B_y) \hat{a}_y + (A_z - B_z) \hat{a}_z\end{aligned}$$

Vektor-scalar multiplication

$$\begin{aligned}a\mathbf{B} &= a(B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= aB_x \hat{a}_x + aB_y \hat{a}_y + aB_z \hat{a}_z \\ &= (aB_x) \hat{a}_x + (aB_y) \hat{a}_y + (aB_z) \hat{a}_z\end{aligned}$$


Dot Product

$$\begin{aligned}\mathbf{A} \cdot \mathbf{B} &= (A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z) \cdot (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x \hat{a}_x \cdot (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &\quad + A_y \hat{a}_y \cdot (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &\quad + A_z \hat{a}_z \cdot (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x B_x (\hat{a}_x \cdot \hat{a}_x) + A_x B_y (\hat{a}_x \cdot \hat{a}_y) + A_x B_z (\hat{a}_x \cdot \hat{a}_z) \\ &\quad + A_y B_x (\hat{a}_y \cdot \hat{a}_x) + A_y B_y (\hat{a}_y \cdot \hat{a}_y) + A_y B_z (\hat{a}_y \cdot \hat{a}_z) \\ &\quad + A_z B_x (\hat{a}_z \cdot \hat{a}_x) + A_z B_y (\hat{a}_z \cdot \hat{a}_y) + A_z B_z (\hat{a}_z \cdot \hat{a}_z) = A_x B_x + A_y B_y + A_z B_z\end{aligned}$$


Cross Product

$$\begin{aligned}\mathbf{A} \times \mathbf{B} &= (A_x \hat{a}_x + A_y \hat{a}_y + A_z \hat{a}_z) \times (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x \hat{a}_x \times (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &\quad + A_y \hat{a}_y \times (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &\quad + A_z \hat{a}_z \times (B_x \hat{a}_x + B_y \hat{a}_y + B_z \hat{a}_z) \\ &= A_x B_x (\hat{a}_x \times \hat{a}_x) + A_x B_y (\hat{a}_x \times \hat{a}_y) + A_x B_z (\hat{a}_x \times \hat{a}_z) \\ &\quad + A_y B_x (\hat{a}_y \times \hat{a}_x) + A_y B_y (\hat{a}_y \times \hat{a}_y) + A_y B_z (\hat{a}_y \times \hat{a}_z) \\ &\quad + A_z B_x (\hat{a}_z \times \hat{a}_x) + A_z B_y (\hat{a}_z \times \hat{a}_y) + A_z B_z (\hat{a}_z \times \hat{a}_z)\end{aligned}$$

$$\mathbf{A} \times \mathbf{B} = (A_y B_z - A_z B_y) \hat{a}_x + (A_z B_x - A_x B_z) \hat{a}_y + (A_x B_y - A_y B_x) \hat{a}_z$$



Coordinate Systems



Latihan soal

1. Diberikan tiga vektor pada sistem koordinat Kartesian dibawah ini :

$$\vec{A} = \hat{a}_x + \hat{a}_y$$

$$\vec{B} = \hat{a}_x + 2\hat{a}_y - 2\hat{a}_z$$

$$\vec{C} = \hat{a}_y + 2\hat{a}_z$$

tentukan hasil dari operasi-operasi vektor dibawah ini :

a) $\vec{A} + \vec{B}$

b) $\vec{B} - \vec{C}$

c) $4\vec{C}$


d) $\vec{A} \bullet \vec{B}$

e) $\vec{A} \times \vec{B}$

f) $\vec{A} \bullet \vec{B} \times \vec{C}$

Coordinate Systems

Elemen Perpindahan, Elemen Luas, dan Elemen Volume




| | | |
|--|---|---|
| Elemen perpindahan $d\vec{L} = dx\hat{a}_x + dy\hat{a}_y + dz\hat{a}_z$ | Elemen perpindahan $d\vec{L} = d\rho \hat{a}_\rho + \rho d\phi \hat{a}_\phi + dz \hat{a}_z$ | Elemen perpindahan $d\vec{L} = dr \hat{a}_r + rd\theta \hat{a}_\theta + r \sin\theta d\phi \hat{a}_\phi$ |
| Elemen luas $d\vec{S}_y = dx dz \hat{a}_y$ $d\vec{S}_x = dy dz \hat{a}_x$ $d\vec{S}_z = dx dy \hat{a}_z$ | Elemen luas $d\vec{S}_\phi = d\rho dz \hat{a}_\phi$ $d\vec{S}_\rho = \rho d\phi dz \hat{a}_\rho$ | Elemen luas $d\vec{S}_\theta = r \sin\theta dr d\phi \hat{a}_\theta$ $d\vec{S}_r = r \sin\theta d\theta d\phi \hat{a}_r$ |
| Elemen volume $dV = dx dy dz$ | Elemen volume $dV = \rho d\rho d\phi dz$ | Elemen volume $dV = r^2 \sin\theta dr d\theta d\phi$ |

Calculus of scalar and Vector



Gradient

- Gradien dari suatu scalar field adalah suatu vektor yang magnitudenya menunjukkan perubahan maksimum scalar field tersebut dan arahnya menunjukkan arah dari peningkatan tercepat scalar field tersebut
- Ilustrasi 1




Suhu adalah scalar field. Jika terukur suhu pada suatu titik X dari sebuah sumber lilin, maka gradien terhadap suhu di X adalah vektor A dan bukan vektor B

- Ilustrasi 2



Ketinggian/kontur permukaan bumi adalah scalar field $h(x,y)$. Jika kita melakukan operasi gradient terhadap $h(x,y)$, maka akan menghasilkan vector field seperti gambar disamping



Calculus of scalar and Vector

Gradient operator pada sistem koordinat

GRADIENT PADA KOORDINAT KARTESIAN

$$\nabla g(\bar{r}) = \frac{\partial g(\bar{r})}{\partial x} \hat{a}_x + \frac{\partial g(\bar{r})}{\partial y} \hat{a}_y + \frac{\partial g(\bar{r})}{\partial z} \hat{a}_z$$

GRADIENT PADA KOORDINAT TABUNG

$$\nabla g(\bar{r}) = \frac{\partial g(\bar{r})}{\partial r} \hat{a}_r + \frac{1}{r} \frac{\partial g(\bar{r})}{\partial \theta} \hat{a}_\theta + \frac{1}{r \sin \theta} \frac{\partial g(\bar{r})}{\partial \phi} \hat{a}_\phi$$

GRADIENT PADA KOORDINAT BOLA

$$\nabla g(\bar{r}) = \frac{\partial g(\bar{r})}{\partial \rho} \hat{a}_\rho + \frac{1}{\rho} \frac{\partial g(\bar{r})}{\partial \phi} \hat{a}_\phi + \frac{\partial g(\bar{r})}{\partial z} \hat{a}_z$$

Contoh

Misalkan,

$$V(x, y, z) = x^2yz^3$$

maka,

$$\vec{E} = -\vec{\nabla}V = -\left[\frac{\partial}{\partial x} \hat{a}_x + \frac{\partial}{\partial y} \hat{a}_y + \frac{\partial}{\partial z} \hat{a}_z \right] [x^2yz^3]$$

$$= -\left[\frac{\partial(x^2yz^3)}{\partial x} \hat{a}_x + \frac{\partial(x^2yz^3)}{\partial y} \hat{a}_y + \frac{\partial(x^2yz^3)}{\partial z} \hat{a}_z \right]$$

$$= -[2xyz^3 \hat{a}_x + x^2z^3 \hat{a}_y + 3x^2yz^2 \hat{a}_z]$$

$$\vec{\nabla} = \frac{\partial}{\partial x} \hat{a}_x + \frac{\partial}{\partial y} \hat{a}_y + \frac{\partial}{\partial z} \hat{a}_z$$

operator Del



Calculus of scalar and Vector


Divergensi

- Untuk mengestimasi dan meng-kuantisasi vector-vector field, sering dengan cara mengukur aliran vector field tersebut (atau netto aliran masuk dan keluar)
- Divergensi mengamati unsur volume tertentu yang sangat kecil, mengamati apakah ada 'sumber' atau tidak di dalam volume tersebut
- Definisi dan simbol: Misalkan vector field yang diamati adalah vektor D

$$\lim_{\Delta V \rightarrow 0} \frac{\int_S \vec{D} \cdot d\vec{S}}{\Delta V}$$

- Hasil operasi divergensi adalah skalar, karena dot product
- Misalkan vector field yang diamati adalah vektor D maka:

- **Hasil divergensi (+)** Jumlah vektor keluar > jumlah vektor masuk
Artinya : Di dalam ruang **ada sumber**
- **Hasil divergensi (-)** Jumlah vektor keluar < jumlah vektor masuk
Artinya : **Ada kekosongan** dalam volume dan bersifat menyerap, contoh : **Black Hole**
- **Hasil divergensi = 0** Jumlah vektor keluar = jumlah vektor masuk
Artinya : **Tidak ada apa-apa** dalam volume tersebut



Calculus of scalar and Vector

Divergensi pada sistem koordinat

DIVERGENSI PADA KOORDINAT KARTESIAN

$$\nabla \cdot \mathbf{A}(\bar{r}) = \frac{\partial A_x(\bar{r})}{\partial x} + \frac{\partial A_y(\bar{r})}{\partial y} + \frac{\partial A_z(\bar{r})}{\partial z}$$

DIVERGENSI PADA KOORDINAT TABUNG

$$\nabla \cdot \mathbf{A}(\bar{r}) = \frac{1}{\rho} \left[\frac{\partial (\rho A_\rho(\bar{r}))}{\partial \rho} \right] + \frac{1}{\rho} \frac{\partial A_\phi(\bar{r})}{\partial \phi} + \frac{\partial A_z(\bar{r})}{\partial z}$$

DIVERGENSI PADA KOORDINAT BOLA

$$\nabla \cdot \mathbf{A}(\bar{r}) = \frac{1}{r^2} \left[\frac{\partial (r^2 A_r(\bar{r}))}{\partial r} \right] + \frac{1}{r \sin \theta} \left[\frac{\partial (\sin \theta A_\theta(\bar{r}))}{\partial \theta} \right] + \frac{1}{r \sin \theta} \frac{\partial A_\phi(\bar{r})}{\partial \phi}$$

Contoh

Misalkan suatu vector Field :

$$\mathbf{A}(\bar{r}) = \frac{\sin \theta}{r} \hat{a}_r$$

Therefore, $A_\theta = 0$ and $A_\phi = 0$, leaving:

$$\begin{aligned}\nabla \cdot \mathbf{A}(\bar{r}) &= \frac{1}{r^2} \left[\frac{\partial (r^2 A_r(\bar{r}))}{\partial r} \right] \\ &= \frac{1}{r^2} \left[\frac{\partial}{\partial r} \left(r^2 \frac{\sin \theta}{r} \right) \right] \\ &= \frac{1}{r^2} \left[\frac{\partial (r \sin \theta)}{\partial r} \right] \\ &= \frac{1}{r^2} [\sin \theta] = \frac{\sin \theta}{r^2}\end{aligned}$$

Calculus of scalar and Vector

Curl (Pusaran)

- Curl adalah ukuran seberapa besar putaran/pusaran/rotasi dari suatu vector field di sekitar titik tertentu
- Rotasi/pusaran terjadi jika adanya ketidakseragaman vector field
- Definisi dan simbol : Misalkan vector field yang diamati adalah vektor \vec{H}


$$\lim_{\Delta S \rightarrow 0} \frac{\oint \vec{H} \cdot d\vec{L}}{\Delta S}$$

simbol

$$\vec{\nabla} \times \vec{H}$$

- Curl adalah integral garis yang membatasi luas yang sangat kecil
- Curl digunakan untuk mengetahui vector field menembus permukaan diferensial yang sangat kecil, yang menyebabkan pusaran medan lain

- Ilustrasi



Rapat arus J yang menembus permukaan dS menimbulkan suatu pusaran/rotasi medan magnetik H

Calculus of scalar and Vector

Curl pada sistem koordinat

CURL PADA KOORDINAT KARTESIAN

$$\nabla \times \mathbf{A}(\bar{r}) = \left[\frac{\partial A_y(\bar{r})}{\partial z} - \frac{\partial A_z(\bar{r})}{\partial y} \right] \hat{a}_x + \left[\frac{\partial A_z(\bar{r})}{\partial x} - \frac{\partial A_x(\bar{r})}{\partial z} \right] \hat{a}_y + \left[\frac{\partial A_x(\bar{r})}{\partial y} - \frac{\partial A_y(\bar{r})}{\partial x} \right] \hat{a}_z$$

CURL PADA KOORDINAT TABUNG

$$\begin{aligned} \nabla \times \mathbf{A}(\bar{r}) = & \left[\frac{1}{\rho} \frac{\partial A_z(\bar{r})}{\partial \phi} - \frac{\partial A_\phi(\bar{r})}{\partial z} \right] \hat{a}_\rho \\ & + \left[\frac{\partial A_\rho(\bar{r})}{\partial z} - \frac{\partial A_z(\bar{r})}{\partial \rho} \right] \hat{a}_\phi \\ & + \left[\frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\phi(\bar{r})) - \frac{1}{\rho} \frac{\partial A_\rho(\bar{r})}{\partial \phi} \right] \hat{a}_z \end{aligned}$$


CURL PADA KOORDINAT BOLA

$$\begin{aligned} \nabla \times \mathbf{A}(\bar{r}) = & \left[\frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta A_\phi(\bar{r})) - \frac{1}{r \sin \theta} \frac{\partial A_\theta(\bar{r})}{\partial \phi} \right] \hat{a}_r \\ & + \left[\frac{1}{r \sin \theta} \frac{\partial A_r(\bar{r})}{\partial \phi} - \frac{1}{r} \frac{\partial}{\partial r} (r A_\phi(\bar{r})) \right] \hat{a}_\theta \\ & + \left[\frac{1}{r} \frac{\partial}{\partial r} (r A_\theta(\bar{r})) - \frac{1}{r} \frac{\partial A_r(\bar{r})}{\partial \theta} \right] \hat{a}_\phi \end{aligned}$$

Divergence Theorem/Gauss's Theorem

□ Teorema Divergensi

$$\iiint_V (\nabla \cdot \vec{F}) dV = \iint_S (\vec{F} \cdot d\vec{S})$$




FLUX: adalah netto aliran yang menembus permukaan dengan arah normal terhadap permukaan


$$\Psi = \iint_S \vec{F} \cdot d\vec{S} = \iint_S F \cos\theta dS$$

$$d\vec{S} = dS \hat{n}$$

Vektor dS selalu tegak lurus terhadap elemen permukaan dS




Teorema divergensi menyatakan bahwa jumlah flux dari suatu vector field yang menembus suatu permukaan tertutup/close surface sama dengan jumlah total / integral volume dari suatu divergensi vector field tersebut pada seluruh area(volume) yang terlingkupi close surface tersebut




Stoke's Theorem

□ Theorema stoke


$$\iint_S \nabla \times \mathbf{A}(\bar{r}) \cdot d\bar{s} = \oint_C \mathbf{A}(\bar{r}) \cdot d\bar{\ell}$$



Contour/Garis tepi C adalah close contour yang mengelilingi permukaan/surface S. arah dari contour C didefinisikan dari vektor ds dan aturan putaran tangan kanan. Pada gambar disamping contour C berputar berlawanan arum jam.



A closed contour adalah contour/garis tepi yang dimulai dan diakhiri pada titik yang sama!



Artinya, jika kita ingin mengintegralkan suatu vector field terhadap suatu open surface maka seolah olah kita ingin menjumlahkan / mengintegralkan suatu pusaran di setiap titik pada suatu permukaan.

$$\iint_S \mathbf{B}(\bar{r}) \cdot d\bar{s} = \iint_S \nabla \times \mathbf{A}(\bar{r}) \cdot d\bar{s}$$

Hal tersebut sama saja dengan secara sederhana menjumlahkan /mengintegralkan 'rotasi' vector field tersebut pada sepanjang close contour yang mengelilingi surface tersebut.

ANY QUESTION???



